

YUMS 97-19
 KEK-TH-526
 DESY 97-131
 August 13, 1997

CP Violation at Electron-Positron Colliders*

M.S. Baek and S.Y. Choi

Department of Physics, Yonsei University, Seoul 120-749, Korea

and

K. Hagiwara

Theory Group, KEK, 1-1 Oho, Tsukuba, Ibaraki 305, Japan

Abstract

A detailed, model-independent study of CP violation at present and future electron-positron colliders is reported. Firstly, we investigate CP violation effects in W boson and t quark pair production in e^+e^- annihilation and in two-photon fusion at a next e^+e^- linear collider, where Compton-backscattered laser light off the electrons or positrons are employed as a powerful polarized photon source. Secondly, noting that there do not exist any direct measurements for the tau-lepton electric dipole moment (EDM), we address the importance of performing its direct measurements at e^+e^- collisions off the Z -boson pole at TRISTAN, LEP II and CLEO II. We present a rough comparison of the potential of these experiments in the τ EDM measurements. Finally, we report on our recent works for probing CP violation in the semileptonic decays of the tau lepton, which involve two different intermediate resonances with large ratios of widths to masses, and which can be most efficiently identified at the planned B and proposed τ -charm factories. All the CP -violation phenomena in the processes under consideration, if discovered, imply new CP -violation mechanisms completely different from the Standard Model CP violation mechanism through the complex phase of the Cabibbo-Kobayashi-Maskawa matrix.

*To appear in the Proceedings of the KEK meetings on ‘ CP violation and its origin’ (1993-1997).

1 Introduction

The Standard Model (SM) of strong and electroweak interactions[1] is widely believed to be the appropriate effective theory below the TeV energy scale. There have been many efforts in testing the SM through precise measurements over a broad range of energies, but there are no confirmed experimental data that disagree with the SM up to now. However, it is believed that the SM is not the final theory of Nature because of several unresolved problems within this model. The SM has too many (at least nineteen) free parameters. This model cannot say anything about the number of families and the origin of Higgs mechanism, which breaks the $SU(2)_L \times U(1)_Y$ electroweak symmetry and is responsible for the masses of weak gauge bosons and fermions. In addition, it does not include gravitational interactions. These motivate a lot of theoretical attempts to understand the SM as a low energy effective theory of a more fundamental theory and several experimental plans to find new phenomena at and beyond the weak scale.

According to the effective field theory[2], effects of more fundamental physics appear as higher dimensional, non-renormalizable operators at the SM scale ($v = 250$ GeV), which are usually suppressed by powers of the characteristic mass ratio between the SM scale and the large mass scale of the fundamental theory. The most efficient way to find the effects of such higher dimensional operators is to investigate the processes or observables which are suppressed in the SM. In this case, dominant contributions can originate from non-renormalizable higher dimensional operators, usually from those with the lowest possible dimension. In this regard, CP -violating processes or observables deserve special attention since the SM contributions to CP -violating processes are often very small.

Ever since CP violation was discovered in the neutral Kaon system in 1964[3], this system has been the only place where CP violation has been observed. The neutral Kaon system CP violation can be understood within the SM, where the CP -violating effects originate from the complex phase of the CKM matrix[4, 5]. Since this phenomenon can also be due to an entirely different source such as a superweak mixing of K^0 and $\bar{K}^0$ [6], there is no complete understanding of the source of the CP violation up to now.

There is another reason to consider CP violation seriously. It is required to explain the baryon asymmetry of the universe[7]. No antimatter is observed in our solar system. Small amount of anti-protons that are observed in high energy cosmic rays are consistent with being secondary products. These are evidence of the baryon asymmetry on the galactic scale. To generate the baryon asymmetry of the universe starting from the symmetric universe, three conditions, first proposed by Sakharov[8], must be satisfied: B (baryon number) violation, C and CP violation, and departure from equilibrium. The need of C and CP violation for the baryogenesis is obvious when we consider the fact that the baryon number is odd under C and CP transformations. It is shown in recent works[9] that the SM CP -violation mechanism is too insufficient to explain the observed baryon asymmetry of the universe.

It is believed that new paradigm of physics should emerge from the more fundamental theory when experiments are able to probe energies of the electroweak scale. The new theory should explain the cosmological baryon asymmetry and may provide us with richer

possibilities for CP violation. In this aspect, the study of CP -violating effects can be an efficient probe of physics beyond the SM.

As an example, the electric dipole moments (EDM) of quarks and leptons are ideal from an experimental point of view for searching evidence of new, non-KM CP violation effects since the KM mechanism itself tends to give negligibly small contributions to such flavor-conserving processes. The SM contribution to the EDM comes from loop corrections. Several works[10] have shown that the EDM of both the neutron and the electron must vanish at the two-loop level. The SM contribution to the EDM appears only at the three-loop level, and hence it is extremely small. For example, the EDM of the electron arising from the CKM mechanism has been estimated to be about $10^{-38} ecm$ [11], which is twelve orders of magnitude below the current limit.

On the other hand, a number of the most plausible and well-motivated extensions of the SM lead to much larger EDM's since these models can give rise to the EDM's at the one-loop level. Furthermore, the EDM of a particle is proportional to some positive powers of its mass because of the spin flip. It is, therefore, naturally expected that the top-quark as the most massive quark, the W boson as the most massive gauge boson and the τ lepton as the most massive lepton are the best candidates to observe non-KM CP violation.

In the present report, we review a series of our works on a detailed, model-independent study of CP violation in processes involving the heavy particles, the W boson, the top quark, and the τ lepton at present and future e^+e^- colliders[12, 13, 14]. In section 2, we investigate CP violation effects in W -boson and t -quark pair production in e^+e^- annihilation and in two-photon fusion at a next linear e^+e^- collider (NLC)[15], where Compton-backscattered laser lights are employed as a powerful polarized photon source[16]. In section 3, by noting that there do not exist direct measurements of the tau-lepton EDM, we address the importance of performing its direct measurements at e^+e^- collisions off the Z boson pole at TRISTAN, LEP II and CLEO II and present a rough comparison of the potential of these experiments in the τ EDM measurements. Also reported in section 3 is our recent works on probing CP violation in the semileptonic τ decays, which involve two different intermediate resonances with large ratios of widths to masses and which can be studied at CLEO II, the planned B factories, and the proposed τ -charm factories[17]. In section 4, we summarize our findings and make a few concluding remarks.

2 CP Violation at Next e^+e^- Linear Colliders

The possibility of having an NLC has increased significantly in the last few years in parallel with a series of international workshops called Physics and Experiments with Linear e^+e^- Colliders[15]. There is general consensus for a 500 GeV NLC with an integrated luminosity of the order of 10 fb^{-1} for the first phase. With such experimental parameters, NLC produces a copious number of W and top-quark pairs. With the expectations that non-KM CP violation effects from new interactions beyond the SM are predominant in the most massive charged gauge boson and fermion systems, the W boson and the t quark,

a lot of works[18, 19, 20, 21, 22, 23, 24, 25, 26, 27] have made important contributions to probing CP violation through the production and decay processes of these two heavy particles in hadron-hadron and e^+e^- collisions. On the other hand, as one important spin-off of NLC highly-energetic photon beams from Compton backscattering of laser light off electrons or positrons can be utilized to realize electron-photon and photon-photon collisions. Therefore, recently non-KM CP violation has been probed in the W -pair and t -pair production processes via two-photon fusion[28, 13, 29].

In this section we review a series of our recent works[12, 13] on a detailed, model-independent study of CP violation in W -boson pair production and top-quark pair production in the e^+e^- and two-photon annihilation at an NLC.

The works prior to our works are classified into two categories according to their emphasized aspects: (i) the classification of spin and angular correlations of the decay products without electron beam polarization[18, 19, 20, 23, 24, 25, 30] and (ii) the use of a few typical CP -odd observables with electron beam polarization[21, 22]. In the first class, they have constructed a tower of CP -odd observables according to their ranks. However, since the W boson and top quark are spin-1 and spin-1/2, the number of CP -odd spin correlations appearing in the processes $e^+e^-(\gamma\gamma) \rightarrow W^+W^-$ and $e^+e^-(\gamma\gamma) \rightarrow t\bar{t}$ is finite so that *all the linearly-independent CP -odd correlations can be completely defined*. Then, all the previously-considered correlations are expressed in terms of the complete set of linearly-independent CP -odd correlations. In the second class, it has been shown that electron beam polarization is very crucial for a few specialized CP -odd correlations. Those works can be easily extended with the complete set of CP -odd correlations in order to investigate which CP -odd correlations depend crucially on electron polarization and which correlations do not.

In order to make our predictions model-independent, we consider all possible leading CP -violating effects which can be induced by any extensions of the SM at the weak interaction scale. To describe CP violation in the W -boson pair production, we follow the effective field theory approach with a linear realization of the symmetry-breaking sector due to some new interactions which involve the Higgs sector and the electroweak gauge bosons. The $SU(2)_L \times U(1)_Y$ electroweak gauge symmetry is imposed in constructing the higher dimensional effective Lagrangian since there is no experimental sign of violation of this symmetry. There exist six relevant dimension-six CP -odd operators. On the other hand, the relevant CP -odd corrections to the top-quark pair production processes are obviously EDM-type couplings of top and anti-top quarks to a photon and Z .

We use two methods in detecting CP violation. One method makes use of the produced W bosons and top-quarks by measuring various spin correlations in their final decay products, and the other method is to employ polarized photon beams to measure various CP -odd polarization asymmetries of the initial states. In the e^+e^- mode, where the initial e^+e^- state is (almost) CP -even due to the very small electron mass[31], only the first method can be used, but in the $\gamma\gamma$ mode both methods can be employed. We concentrate on the second method in the two-photon mode by making use of the Compton backscattered laser light off the electron or positron beam as a powerful photon source. The polarization of the scattered high energy photon beams can be controlled by adjusting

polarizations of initial electron beams and the laser light. Nevertheless, we will simply consider purely linearly-polarized photon beams in the present report.

2.1 Photon Spectrum

Generally, a purely polarized photon beam state is a linear combination of two helicity states and the photon polarization vector can be expressed in terms of two angles α and ϕ in a given coordinate system as

$$|\alpha, \phi\rangle = -\cos(\alpha)e^{-i\phi}|+\rangle + \sin(\alpha)e^{i\phi}|-\rangle, \quad (1)$$

where $0 \leq \alpha \leq \pi/2$ and $0 \leq \phi \leq 2\pi$. The degrees of circular and linear polarization are $\xi = \cos(2\alpha)$ and $\eta = \sin(2\alpha)$, respectively, and the direction of maximal linear polarization is denoted by the azimuthal angle ϕ . For a partially polarized photon beam it is necessary to re-scale ξ and η by its degree of polarization.




Figure 1: The coordinate system in the colliding $\gamma\gamma$ c.m. frame. The scattering angle, Θ , and the azimuthal angles, ϕ_1 and ϕ_2 , for the linear polarization directions measured from the scattering plane are described.

The state vector of the two-photon system in the c.m. frame, where one photon momentum is along the positive z direction, is

$$|\alpha_1, \phi_1; \alpha_2, \phi_2\rangle = |\alpha_1, \phi_1\rangle |\alpha_2, -\phi_2\rangle. \quad (2)$$

The angle ϕ_1 (ϕ_2) is the azimuthal angle of the maximal linear polarization of the photon beam, whose momentum is in the positive (negative) z direction, with respect to the direction of the $W^+(t)$ momentum in $\gamma\gamma \rightarrow W^+W^- (\gamma\gamma \rightarrow t\bar{t})$ as shown in Fig. 1.

Generally, the angular dependence for $\gamma\gamma \rightarrow X$ with two linearly-polarized photon beams, of which the degrees of linear polarization are η and $\bar{\eta}$, is expressed in the form

$$\mathcal{D}(\eta, \bar{\eta}; \chi, \phi) = \Sigma_{\text{unpol}} - \frac{1}{2} \text{Re} \left[\left(\eta e^{-i\phi} + \bar{\eta} e^{i\phi} \right) e^{-i\chi} \Sigma_{02} \right]$$

$$+\frac{1}{2}Re\left[\left(\eta e^{-i\phi}-\bar{\eta}e^{i\phi}\right)e^{-i\chi}\Delta_{02}\right]+\eta\bar{\eta}Re\left[e^{-2i\phi}\Sigma_{22}+e^{-2i\chi}\Sigma_{00}\right], \quad (3)$$

with the subscripts, 0 and 2, representing the magnitude of the sum of two photon helicities of the initial two-photon system. Here, two angular variables χ and ϕ are given by

$$\chi = \phi_1 - \phi_2, \quad \phi = \phi_1 + \phi_2. \quad (4)$$

The azimuthal angle difference, χ , is independent of the final state, while the azimuthal angle sum, ϕ , depends on the scattering plane, and both angles are invariant with respect to the Lorentz boost along the two-photon beam direction.

Table 1: CP and CPT properties of the invariant functions and the angular distributions.

CP	CPT	Invariant functions	Angular dependences
even	even	Σ_{unpol}	
		$\mathcal{R}(\Sigma_{02})$	$\eta \cos(\phi + \chi) + \bar{\eta} \cos(\phi - \chi)$
		$\mathcal{R}(\Sigma_{22})$	$\eta\bar{\eta} \cos(2\phi)$
		$\mathcal{R}(\Sigma_{00})$	$\eta\bar{\eta} \cos(2\chi)$
even	odd	$\mathcal{I}(\Delta_{02})$	$\eta \sin(\phi + \chi) + \bar{\eta} \sin(\phi - \chi)$
		$\mathcal{I}(\Sigma_{22})$	$\eta\bar{\eta} \sin(2\phi)$
odd	even	$\mathcal{I}(\Sigma_{02})$	$\eta \sin(\phi + \chi) - \bar{\eta} \sin(\phi - \chi)$
		$\mathcal{I}(\Sigma_{00})$	$\eta\bar{\eta} \sin(2\chi)$
odd	odd	$\mathcal{R}(\Delta_{02})$	$\eta \cos(\phi + \chi) - \bar{\eta} \cos(\phi - \chi)$

It is useful to classify the invariant functions Σ and Δ 's, which depend only on the scattering angle Θ , and the angular distributions according to their transformation properties under the discrete symmetries, CP and CPT ^a as shown in Table 1. There exist three CP -odd functions; $\mathcal{I}(\Sigma_{02})$, $\mathcal{I}(\Sigma_{00})$ and $\mathcal{R}(\Delta_{02})$. Here, $\mathcal{R}$ and $\mathcal{I}$ stand for real and imaginary parts, respectively. While the first two terms are CPT -even, the last term $\mathcal{R}(\Delta_{02})$ is CPT -odd. Since the CPT -odd term $\mathcal{R}(\Delta_{02})$ requires an absorptive part in the amplitude, it is generally expected to be smaller in magnitude than the CPT -even terms. We then can define two CP -odd asymmetries from $\mathcal{I}(\Sigma_{02})$ and $\mathcal{I}(\Sigma_{00})$. First, note that Σ_{00} is independent of the azimuthal angle ϕ whereas Σ_{02} is not. In order to improve the observability we may integrate $\mathcal{I}(\Sigma_{02})$ over the azimuthal angle ϕ with an appropriate weight function. Without any loss of generality we can take $\eta = \bar{\eta}$. Then, $\mathcal{I}(\Sigma_{00})$ can be separated by taking the difference of the distributions at $\chi = \pm\pi/4$ and the $\mathcal{I}(\Sigma_{02})$ by taking the difference of the distributions at $\chi = \pm\pi/2$. As a result we obtain the following

^a $\tilde{T}$ is the "naive" time reversal operation which flips particle momenta and spins, but does not interchange initial and final states

two integrated CP -odd asymmetries^b:

$$\hat{A}_{02} = \left(\frac{2}{\pi}\right) \frac{\mathcal{I}(\Sigma_{02})}{\Sigma_{\text{unpol}}}, \quad \hat{A}_{00} = \frac{\mathcal{I}(\Sigma_{00})}{\Sigma_{\text{unpol}}}, \quad (5)$$

where the factor $(2/\pi)$ in the $\hat{A}_{02}$ stems from taking the average over the azimuthal angle ϕ with the weight function $\text{sign}(\cos \phi)$.

The Compton backscattering process is characterized by two parameters x and y :

$$x = \frac{4E\omega_0}{m_e^2} \approx 15.3 \left(\frac{E}{\text{TeV}}\right) \left(\frac{\omega_0}{\text{eV}}\right), \quad y = \frac{\omega}{E}, \quad (6)$$

where E is the electron beam energy and ω_0 the incident laser beam frequency. On the average, the backscattered photon energies increase with x ; the maximum photon energy fraction is given by $y_m = x/(1+x)$. Operation below the threshold[16] for e^+e^- pair production in collisions between the laser beam and the Compton-backscattered photon beam requires $x \leq 2(1 + \sqrt{2}) \approx 4.83$.

In the two-photon collision case with Compton-backscattered photon beams, only part of each laser linear polarization is transferred to the high-energy photon beam and the degrees of linear polarization transfer^c are determined by two functions, $\mathcal{A}_\eta$ and $\mathcal{A}_{\eta\eta}$:

$$\mathcal{A}_\eta(\tau) = \frac{\langle \phi_0 \phi_3 \rangle_\tau}{\langle \phi_0 \phi_0 \rangle_\tau}, \quad \mathcal{A}_{\eta\eta}(\tau) = \frac{\langle \phi_3 \phi_3 \rangle_\tau}{\langle \phi_0 \phi_0 \rangle_\tau}, \quad (7)$$

where $\phi_0(y) = \frac{1}{1-y} + 1 - y - 4r(1-r)$ and $\phi_3(y) = 2r^2$ with $r = y/x(1-y)$, and τ is the ratio of the $\gamma\gamma$ c.m. energy squared $\hat{s}$ to the e^+e^- collider energy squared s . $\mathcal{A}_\eta$ is for the collision of an unpolarized photon beam and a linearly polarized photon beam, and $\mathcal{A}_{\eta\eta}$ for that of two linearly polarized photon beams. $\langle \phi_i \phi_j \rangle_\tau$ ($i, j = 0, 3$) is defined as a normalized convolution integral for a fixed value of τ . Folding the photon spectrum with the $\gamma\gamma \rightarrow X$ cross section yields two CP -odd asymmetries A_{02} and A_{00} , which depend crucially on the two-photon spectrum and the two linear polarization transfers.

We first investigate the $\sqrt{\tau}$ dependence of the two-photon spectrum and the two linear polarization transfers, $\mathcal{A}_\eta$ and $\mathcal{A}_{\eta\eta}$ by varying the value of the parameter x . Three values of x are chosen; $x = 0.5, 1$, and 4.83 . Fig. 2 clearly shows that the energy of two photons reaches higher ends for larger x values but the maximum linear polarization transfers are larger for smaller x values. The parameter x should be kept as large as possible to reach higher energies. However, larger CP -odd asymmetries can be obtained for smaller x values. Therefore, there should exist a compromise value of x for the optimal observability of CP violation. In this regard, it is very crucial to have a laser with adjustable beam frequency.

^bThe authors in Ref. [29] have also considered $\mathcal{I}(\Sigma_{02})$, but not $\mathcal{I}(\Sigma_{00})$

^cThe function $A_{\eta\eta}$ has been considered in Ref. [32].




Figure 2: (a) the $\gamma\gamma$ luminosity spectrum and (b) the two linear polarization transfers, $\mathcal{A}_\eta$ (solid lines) and $\mathcal{A}_{\eta\eta}$ (dashed lines), for $x = 4E\omega_0/m_e^2 = 0.5, 1$ and 4.83 .

2.2 W pair production

To describe CP violation from new interactions among electroweak vector bosons in a model-independent way[33], we assume that the operators with lowest energy dimension dominate the CP -odd amplitudes and that they respect the electroweak gauge invariance which is broken spontaneously by an effective $SU(2)_L$ -doublet scalar. Considering CP -odd interactions of dimension six composed of electroweak gauge bosons and Higgs fields, we can construct an effective Lagrangian $\mathcal{L}_{\text{eff}}$, which is the sum of the SM Lagrangian $\mathcal{L}_{\text{SM}}$ and the new Lagrangian $\mathcal{L}_{\text{new}}$ defined as

$$\mathcal{L}_{\text{new}} = \frac{1}{\Lambda^2} \left[f_{B\tilde{B}} \mathcal{O}_{B\tilde{B}} + f_{B\tilde{W}} \mathcal{O}_{B\tilde{W}} + f_{W\tilde{W}} \mathcal{O}_{W\tilde{W}} + f_{\tilde{B}} \mathcal{O}_{\tilde{B}} + f_{\tilde{W}} \mathcal{O}_{\tilde{W}} + f_{WW\tilde{W}} \mathcal{O}_{WW\tilde{W}} \right], \quad (8)$$

where the six CP -odd dimension-six operators are as follows

$$\begin{aligned} \mathcal{O}_{B\tilde{B}} &= g'^2 (\Phi^\dagger \Phi) B_{\mu\nu} \tilde{B}^{\mu\nu}, & \mathcal{O}_{B\tilde{W}} &= gg' (\Phi^\dagger \sigma^I \Phi) B_{\mu\nu} \tilde{W}^{I\mu\nu}, \\ \mathcal{O}_{W\tilde{W}} &= g^2 (\Phi^\dagger \Phi) W_{\mu\nu}^I \tilde{W}^{I\mu\nu}, & \mathcal{O}_{\tilde{B}} &= ig' [(D_\mu \Phi)^\dagger (D_\nu \Phi)] \tilde{B}^{\mu\nu}, \\ \mathcal{O}_{\tilde{W}} &= ig [(D_\mu \Phi)^\dagger \sigma^I (D_\nu \Phi)] \tilde{W}^{I\mu\nu}, & \mathcal{O}_{WW\tilde{W}} &= g^3 \epsilon^{IJK} \tilde{W}^{I\mu\nu} W_\nu^{J\rho} W_{\rho\mu}^K, \end{aligned} \quad (9)$$

with the Higgs-doublet Φ , $\tilde{W}^{I\mu\nu} = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} W_{\alpha\beta}^I$, and $\tilde{B}^{\mu\nu} = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} \tilde{B}_{\alpha\beta}$. Table 2 shows which vertices already exist in the SM at tree level and which new vertices appear from the new dimension-six CP -odd operators.

For the sake of an efficient analysis in the following, we define four new dimensionless form factors, Y_i ($i = 1$ to 4), which are related with the coefficients, f_i 's ($i = B\tilde{B}, B\tilde{W}, W\tilde{W}, \tilde{B}, \tilde{W}, WW\tilde{W}$) as

$$\begin{aligned} Y_1 &= \left(\frac{m_W}{\Lambda} \right)^2 \left[f_{B\tilde{W}} + \frac{1}{4} f_{\tilde{B}} + f_{\tilde{W}} \right], & Y_2 &= \left(\frac{m_W}{\Lambda} \right)^2 \frac{g^2}{4} f_{WW\tilde{W}}, \\ Y_3 &= \left(\frac{m_W}{\Lambda} \right)^2 \left[f_{W\tilde{W}} + \frac{1}{4} f_{\tilde{W}} \right], & Y_4 &= \left(\frac{m_W}{\Lambda} \right)^2 \left[f_{B\tilde{B}} - f_{B\tilde{W}} - f_{W\tilde{W}} \right]. \end{aligned} \quad (10)$$

Table 2: Vertices relevant for the processes $e^+e^- \rightarrow W^+W^-$ and $\gamma\gamma \rightarrow W^+W^-$ in the effective Lagrangian with the six dimension-six CP -odd operators.

Vertex	γWW	ZWW	$\gamma\gamma WW$	HWW	$\gamma\gamma H$
SM	O	O	O	O	X
$\mathcal{O}_{B\tilde{B}}$	X	X	X	X	O
$\mathcal{O}_{B\tilde{W}}$	O	O	X	X	O
$\mathcal{O}_{W\tilde{W}}$	X	X	X	O	O
$\mathcal{O}_{\tilde{B}}$	O	O	X	X	X
$\mathcal{O}_{\tilde{W}}$	O	O	X	O	X
$\mathcal{O}_{WW\tilde{W}}$	O	O	O	X	X

We note in passing that if all the coefficients, f_i , are of the similar size, then Y_2 would be about ten times smaller than the other form factors in size because of the factor $g^2/4 \sim 0.1$.

2.2.1 Electron-positron mode

The angular dependence $\Sigma_{L,R}(\Theta; \theta, \bar{\theta}; \phi, \bar{\phi})$ of a sequential process $e^+e^- \rightarrow W^+W^- \rightarrow (f_1\bar{f}_2)(f_3\bar{f}_4)$ can be decomposed in terms of eighty-one orthogonal functions $\mathcal{D}_\alpha$'s of the W^+ and W^- decay products as

$$\Sigma_{L,R}(\Theta; \theta, \bar{\theta}; \phi, \bar{\phi}) = \sum_{\alpha=1}^{81} \mathcal{P}_{\alpha L,R}(\Theta) \mathcal{D}_\alpha(\theta, \bar{\theta}; \phi, \bar{\phi}), \quad (11)$$

where Θ is the scattering angle between e^- and W^- , and $\theta(\bar{\theta})$ and $\phi(\bar{\phi})$ are the angular variables of the W^- and W^+ decay products. All the terms, $\mathcal{P}_{\alpha X}$ and $\mathcal{D}_\alpha$, can be divided into four categories under CP and CPT : even-even, even-odd, odd-even, and odd-odd terms[12]. There exist *thirty-six* independent CP -odd terms among which *eighteen* terms are CPT -even and the other *eighteen* CPT -odd.

Including electron polarization, most of the works[23, 24, 20, 25] prior to our works have considered four special CP -odd and CPT -even asymmetries, of which two are essentially equivalent to the so-called triple vector products, and four new CP -odd and CPT -odd asymmetries in addition to the two conventional lepton energy asymmetries. Clearly, the present analysis shows that much more CP -odd asymmetries are available.

Observables which are constructed from the momenta of the charged leptons originating from W^+ and W^- decay are directly and most easily measurable in future experiments. Therefore, the W leptonic decay channels together with the corresponding charge-conjugated ones are exclusively used in the following analysis. The first set of observables under consideration involves the momentum of a lepton from W^- decay correlated with the momentum of a lepton from W^+ decay in the sequential process

$$e^+(\vec{p}_e) + e^-(\vec{p}_e) \rightarrow W^+ + W^- \rightarrow l^+(\vec{q}_+) + l^-(\vec{q}_-) + X, \quad (12)$$

As shown in Ref. [34] a tower of CP -odd observables can be in principle constructed, among which a few typical CP -odd observables[34] are listed in the following:

$$\begin{aligned}
A_1 &= \hat{p}_e \cdot (\vec{q}_+ \times \vec{q}_-), \\
T_{ij} &= (\vec{q}_- - \vec{q}_+)_i (\vec{q}_- \times \vec{q}_+)_j + (i \leftrightarrow j), \\
A_E &= E_+ - E_-, \quad A_2 = \hat{p}_e \cdot (\vec{q}_+ + \vec{q}_-), \\
Q_{ij} &= (\vec{q}_- + \vec{q}_+)_i (\vec{q}_- - \vec{q}_+)_j + (i \leftrightarrow j) - \frac{2}{3} \delta_{ij} (\vec{q}_-^2 - \vec{q}_+^2).
\end{aligned} \tag{13}$$

The observables T and A_1 are CPT -even, whereas Q , A_2 and A_E are CPT -odd. Certainly, all the CP -odd observables can be expressed as a linear combination of a fixed number of linearly-independent observables whose classification depends only on the spins of the final particles[12]. Because of the lack of space, we refer to the work[12] for a more detailed explanation for the point and, following the same procedure as Ref. [34], we use the observables (13) in probing CP violation in $e^+e^- \rightarrow W^+W^-$ in the present report.

The statistical significance of a given observable O_X is determined by comparing its expectation $\langle O_X \rangle$ with the expectation variance $\langle O_X^2 \rangle_{\text{SM}}$ in the SM where $\langle O_X \rangle_{\text{SM}}$ vanishes. Quantitatively, an observation of any deviation from the SM expectation with better than one-standard deviations requires

$$\langle O_X \rangle \geq \sqrt{\frac{\langle O_X^2 \rangle_{\text{SM}}}{N_{WW}}}, \quad N_{WW} = \varepsilon [B_{X^+} \bar{B}_{X^-}] \mathcal{L}_{ee} \sigma(e^+e^- \rightarrow W^+W^-), \tag{14}$$

where N_{WW} is the number of events, $\mathcal{L}_{ee}$ is the e^+e^- integrated luminosity, $B_{X^\pm}$ are the branching fractions of $W^\pm \rightarrow l^\pm \nu_l$, and ε is the detection efficiency, which is assumed to be unity.




Figure 3: (a) The $1\text{-}\sigma$ allowed regions of $Re(Y_1)$ and $Re(Y_2)$ through A_1^l (solid) and T_{33}^l (long-dashed) with polarized electron beams and with the e^+e^- integrated luminosity 10 fb^{-1} at $\sqrt{s} = 500 \text{ GeV}$. (b) The $1\text{-}\sigma$ allowed regions of $Im(Y_1)$ and $Im(Y_2)$ through A_E^l (solid), A_2^l (long-dashed) and Q_{33}^l (dashed) with polarized electron beams and with the e^+e^- integrated luminosity 10 fb^{-1} at $\sqrt{s} = 500 \text{ GeV}$.

In determining the $1\text{-}\sigma$ allowed region of the (Y_1, Y_2) plane we take the following set of experimental parameters:

$$\sqrt{s} = 0.5 \text{ TeV}, \quad L_{ee} = 10 \text{ fb}^{-1}. \quad (15)$$

We note in Fig. 3 that the use of longitudinal electron beam polarization obviates the need for the simultaneous measurement of more than one distribution and it can greatly enhance the sensitivities to the CP -odd parameters by using two or more CP -odd observables. The $1\text{-}\sigma$ optimal sensitivities to $(\text{Re}(Y_1), \text{Re}(Y_2))$ and $(\text{Im}(Y_1), \text{Im}(Y_2))$ read

$$\begin{aligned} |\text{Re}(Y_1)| &\leq 2.9 \times 10^{-3}, & |\text{Re}(Y_2)| &\leq 3.7 \times 10^{-4}, \\ |\text{Im}(Y_1)| &\leq 3.9 \times 10^{-3}, & |\text{Im}(Y_2)| &\leq 1.0 \times 10^{-3}. \end{aligned} \quad (16)$$

2.2.2 Two-photon mode

In counting experiments where final W polarizations are not analyzed, we measure only the distributions summed over the final W polarizations, from which the explicit form of Σ_{unpol} , Σ_{02} , Δ_{02} , Σ_{22} , and Σ_{00} in Eq. (3) can be obtained. First of all, we emphasize that $\mathcal{I}(\Sigma_{00})$ does not require any identification of the scattering plane as mentioned before. Even if one excludes the $\tau^+\tau^- + \not{p}$ modes of 1%, the remaining 99% of the events can be used to measure $\mathcal{I}(\Sigma_{00})$. On the other hand, the scattering plane should be identified to measure $\mathcal{I}(\Sigma_{02})$. Nevertheless, it is worth noting that the charge of the decaying W is not needed to extract $\mathcal{I}(\Sigma_{02})$. Therefore, all the modes except for the $l^+l^- + \not{p}$ modes (9%) can be used for $\mathcal{I}(\Sigma_{02})$.

The $\gamma\gamma \rightarrow W^+W^-$ reaction has a much larger cross section than heavy fermion-pair production such as $\gamma\gamma \rightarrow t\bar{t}$ and, furthermore, the total cross section approaches a constant value at high energies. At $\sqrt{s} = 500 \text{ GeV}$ the total cross section is about 80 pb, while the $t\bar{t}$ cross section is about 1 pb. So, there exist no severe background problems. In the following analysis we simply assume that all the W pair events can be used.

We present our numerical results for the following set of experimental parameters:

$$\sqrt{s} = 0.5 \text{ and } 1.0 \text{ TeV}, \quad \kappa^2 L_{ee} = 20 \text{ fb}^{-1}. \quad (17)$$

Here, κ is the $e\text{-}\gamma$ conversion coefficient in the Compton backscattering. The parameter x , which is dependent on the laser frequency ω_0 , is treated as an adjustable parameter. Folding the photon luminosity spectrum and integrating the distributions over the final kinematic variables, we obtain the x -dependence of available event rates.

Separating the CP -odd asymmetries A_a into three parts as

$$A_a = R(Y_1)A_a^{Y_1} + R(Y_2)A_a^{Y_2} + R(Y_4)A_a^{Y_4}, \quad (18)$$

and considering each form factor separately, we then obtain the $1\text{-}\sigma$ allowed upper bounds of the form factors ($i = 1, 2, 4$)

$$\text{Max}(|R(Y_i)|_a) = \frac{\sqrt{2}}{|A_a^{Y_i}| \sqrt{\varepsilon N_{\text{unpol}}}}, \quad (19)$$




Figure 4: The x dependence of $\text{Max}(|R(Y_1)|)$ and $\text{Max}(|R(Y_2)|)$ at $\sqrt{s} = 0.5$ (solid) and 1.0 TeV (long-dashed) from (a) A_{02} and (b) A_{00} .

if no asymmetry is found. Here, ε is for the sum of W branching fractions available, which is taken to be 100% for A_{00} and 91% for A_{02} .

Postponing the analysis of the CP -violation effects due to Y_4 [12, 26], we present the analysis of the constraints on Y_1 and Y_2 . Figs. 4(a) and (b) show the x dependence of the 1- σ sensitivities to $R(Y_1)$ and $R(Y_2)$, which are obtained from A_{02} and A_{00} , respectively, for $\sqrt{s} = 0.5$ TeV (solid) and $\sqrt{s} = 1$ TeV (long-dashed). These figures and Table 3 clearly exhibit that (i) the sensitivities, especially through A_{00} , depend strongly on x , and (ii) the optimal sensitivities on $R(Y_2)$ are very much improved as $\sqrt{s}$ increases from 0.5 TeV to 1 TeV, while those of $R(Y_1)$ are a little improved.

Table 3: The best 1- σ sensitivities to $R(Y_1)$ and $R(Y_2)$, and their corresponding x values for $\sqrt{s} = 0.5$ and 1 TeV.

Asymmetry	A_{02}		A_{00}	
$\sqrt{s}$ (TeV)	0.5	1.0	0.5	1.0
x	1.83	0.96	0.75	0.31
$\text{Max}(R(Y_1))$	1.1×10^{-2}	5.0×10^{-3}	3.2×10^{-3}	2.2×10^{-3}
x	2.09	1.23	1.11	0.59
$\text{Max}(R(Y_2))$	2.4×10^{-4}	9.0×10^{-5}	2.6×10^{-4}	1.1×10^{-4}

2.3 Top-quark pair production

An important property of a heavy top[35, 36] is that it decays before it can form hadronic bound states[37]. This implies in particular that spin effects, for instance polarization of, and spin correlations between t and $\bar{t}$ quarks can be analyzed through the distributions and angular correlations of the weak decay products of the t and $\bar{t}$ quarks. Moreover, these effects can be calculated in perturbation theory. They provide an additional means for testing SM predictions and of searching for possible new physics effects in top quark production and decay.

The $\gamma t\bar{t}$ vertex consists of not only the SM tree-level vector and axial-vector coupling terms but also a magnetic dipole moment (MDM) and an EDM coupling. Likewise, in addition to the tree-level SM $Zt\bar{t}$ coupling, we have the analogous Z MDM and Z EDM couplings, of which the latter is called the top-quark weak dipole moment (WDM). The MDM-like couplings are present in the SM at the one-loop level. On the other hand, the EDM-like couplings violate CP and, due to the structure of the SM, are only present perturbatively in the SM at the three loop level[10]. In some extensions to the SM such as left-right models, multi-Higgs-doublet models (MHD), supersymmetric SM, however, the EDM couplings may be present even at one-loop level[38, 39]. Neglecting the MDM couplings, we assume the γtt and Ztt vertices to be given by

$$\Gamma_\mu^V = v_V \gamma_\mu + a_V \gamma_\mu \gamma_5 + \frac{c_V}{2m_t} \sigma_{\mu\nu} \gamma_5 q^\nu, \quad V = \gamma, Z, \quad (20)$$

with the vector and axial-vector couplings of the top-quark given in the SM. Here, q is the four-momentum of the vector boson, $V(= \gamma, Z)$. Then, for $m_t = 175$ GeV, the top-quark EDM and WDM, $d_{\gamma,Z}$, are related with $c_{\gamma,Z}$ as

$$d_{\gamma,Z} = \frac{e}{m_t} c_{\gamma,Z} \approx 1.13 \times 10^{-16} c_{\gamma,Z} (ecm). \quad (21)$$

2.3.1 Electron-positron mode

The angular dependence for the process $e^+e^- \rightarrow t\bar{t} \rightarrow (X^+b)(X^-\bar{b})$, can be written as

$$\Sigma_{L,R}(\Theta; \theta, \bar{\theta}; \phi, \bar{\phi}) = \sum_{i=1}^{16} \mathcal{P}_{iL,R}(\Theta) \mathcal{D}_i(\theta, \bar{\theta}; \phi, \bar{\phi}). \quad (22)$$

where Θ is the scattering angle for $e^+e^- \rightarrow t\bar{t}$, and $\theta(\bar{\theta})$ and $\phi(\bar{\phi})$ are the angle variables for the $b(\bar{b})$ in the inclusive decays and for the $l^+(l^-)$ in the semileptonic decays[40] of the top and anti-top quarks, respectively. The terms, $\mathcal{P}_\alpha$ and $\mathcal{D}_\alpha$, can thus be divided into four categories under CP and CPT [13]. There exist *six* independent CP -odd terms among which *three* terms are CPT -even, and the other *three* CPT -odd.

Including electron beam polarization, Poulose and Rindani[22] recently have considered two new CP -odd and CPT -even asymmetries, of which one asymmetry is essentially equivalent to the so-called triple vector product, and two new CP -odd and CPT -odd asymmetries in addition to the two conventional lepton energy asymmetries. Clearly,




Figure 5: The $1\text{-}\sigma$ allowed region of $\text{Re}(c_\gamma)$ and $\text{Re}(c_Z)$ through (a) A_1^b (solid) and T_{33}^b (long-dashed) and (b) A_1^l (solid) and T_{33}^l (long-dashed) with the e^+e^- integrated luminosity 10 fb^{-1} for the polarized electron beam (upper part) and 20 fb^{-1} for the unpolarized electron beam (lower part), respectively, at $\sqrt{s} = 500\text{ GeV}$.

that we can use *six* more asymmetries among which *four* asymmetries are CP -odd and CPT -even and the other *two* terms are CP -odd and CPT -odd.

Observables which are constructed from the momenta of the charged leptons and/or b jets originating from t and $\bar{t}$ decay are directly measurable in future experiments. We consider both the inclusive and exclusive semileptonic decays

$$t \rightarrow bX_{\text{had}}, \quad t \rightarrow bl^+\nu; \quad l = e, \mu, \tau, \quad (23)$$

together with the corresponding charge-conjugated ones, and we use the CPT -even T_{33} and A_1 and the CPT -odd Q_{33} , A_2 and A_E as in the W -pair production case. All the observables are constructed from the momenta of the final b and $\bar{b}$ in the inclusive decay mode and of the final l^+ and l^- in the exclusive semileptonic decay mode.

Inserting the values of the SM electron vector and axial-vector couplings, we obtain for left-handed and right-handed electrons

$$c_L = c_\gamma + 0.64\delta_Z c_Z, \quad c_R = c_\gamma - 0.55\delta_Z c_Z, \quad (24)$$

where $\delta_Z = (1 - m_Z^2/s)^{-1}$. For $m_t = 175\text{ GeV}$ and $m_Z = 91.2\text{ GeV}$, $1 \leq \delta_Z \leq 1.073$. The c_Z contribution to c_L and c_R is similar in size but different in sign. Naturally, electron polarization plays a crucial role in discriminating c_γ and c_Z .

Our numerical results are presented for the following set of experimental parameters:

$$\sqrt{s} = 0.5\text{ TeV}, \quad L_{ee} = \begin{cases} 20\text{ fb}^{-1} & \text{for unpolarized electrons} \\ 10\text{ fb}^{-1} & \text{for polarized electrons} \end{cases}, \quad (25)$$




Figure 6: The $1\text{-}\sigma$ allowed region of $Im(c_\gamma)$ and $Im(c_Z)$ through (a) A_E^b (solid), A_2^b (long-dashed) and Q_{33}^b (dashed) and (b) A_E^l (solid), A_2^l (long-dashed) and Q_{33}^l (dashed) with the e^+e^- integrated luminosity 10 fb^{-1} for the polarized electron beam (upper part) and 20 fb^{-1} for the unpolarized electron beam (lower part) at $\sqrt{s} = 500\text{ GeV}$.

The shadowed parts in Fig 5 show the $1\text{-}\sigma$ allowed regions of $Re(c_\gamma)$ and $Re(c_Z)$ through (a) A_1^b and T_{33}^b and (b) A_1^l and T_{33}^l with polarized electron beams (upper part), respectively, and those in Fig. 6 show the $1\text{-}\sigma$ allowed regions of $Im(c_\gamma)$ and $Im(c_Z)$ through (a) A_E^b , A_2^b and Q_{33}^b and (b) A_E^l , A_2^l and Q_{33}^l with unpolarized electron beams (lower part), respectively. We observe several interesting properties from the figures: (i) The allowed regions strongly depend on electron polarization. (ii) Even with unpolarized electrons and positrons, it is possible to obtain a closed region for the CP -odd parameters by using two or more CP -odd asymmetries. (iii) With polarized electrons, the most stringent bounds on the CPT -even and CPT -odd parameters are obtained through A_1^b and A_E^b in the inclusive top-quark decay mode, respectively. Numerically, the $1\text{-}\sigma$ allowed regions of $Re(c_\gamma)$, $Re(c_Z)$, $Im(c_\gamma)$, and $Im(c_Z)$ are

$$|Re(c_\gamma)| \leq 0.12, \quad |Re(c_Z)| \leq 0.20, \quad (26)$$

$$|Im(c_\gamma)| \leq 0.16, \quad |Im(c_Z)| \leq 0.27. \quad (27)$$

2.3.2 Two-photon mode

Extracting $\mathcal{I}(\Sigma_{00})$ in $\gamma\gamma \rightarrow t\bar{t}$ again does not require identifying the scattering plane, and although the $\tau^+\tau^- + \cancel{p}$ modes of 1% is excluded, the remaining 99% of the events can be used to measure $\mathcal{I}(\Sigma_{00})$. On the other hand, $\mathcal{I}(\Sigma_{02})$ turns out to be zero so that it is useless in determining $Re(c_\gamma)$.

We present our numerical results for the same set of experimental parameters as the process $\gamma\gamma \rightarrow W^+W^-$. Experimentally, $\gamma\gamma \rightarrow W^+W^-$ is the most severe background




Figure 7: The x dependence of the $Re(c_\gamma)$ upper bound, $\text{Max}(|Re(c_\gamma)|)$, at $\sqrt{s} = 0.5$ (solid) and 1 TeV (long-dashed), from A_{00} .

process against $\gamma\gamma \rightarrow t\bar{t}$. In our analysis, we simply take for a numerical analysis a rather conservative value of the detection efficiency $\varepsilon = 10\%$, even though more experimental analysis is required to estimate the efficiency precisely.

It is clear from Fig. 7 and Table 4 that the constraints on $Re(c_\gamma)$ through A_{00} are very sensitive to x and $\sqrt{s}$. It is impressive that the doubling of the c.m. energy enables us to improve the sensitivities almost by a factor of ten.

Table 4: The optimal 1- σ sensitivities to $Re(c_\gamma)$ and their corresponding x values for $\sqrt{s} = 0.5$ and 1 TeV.

$\sqrt{s}$ (TeV)	0.5	1.0
x	3.43	0.85
$Re(c_\gamma)$	0.16	0.02

3 CP Violation in the Tau Lepton System

The τ has the same interaction structure as the e and μ in the SM, apart from their masses. However for practical purposes[17] the τ lepton, the most massive of the known leptons, behaves quite differently from the e and μ leptons in that (i) the τ has hadronic decay modes (e.g. $\tau \rightarrow \pi\nu, \rho\nu, a_1\nu, K^*, \dots$) which allow an efficient measurement of its polarization[41] and (ii) the coupling to the neutral and charged Higgs bosons[42, 43] and other scalar particles is expected to dominate those of the e and μ . These features allow the τ to be a rather special experimental probe of new physics[44, 45, 46, 47, 48, 49].

3.1 Tau Lepton EDM

Recently, the OPAL[50] and ALEPH[51] detector groups have demonstrated by detailed investigations of $Z \rightarrow \tau^+\tau^-$ at LEPI that sensitive CP symmetry tests for the τ WDM at

the few per mill level can be performed in high energy e^+e^- collisions. The OPAL group has employed two optimal genuine CP -odd observables, of which one is CPT -even and the other one CPT -odd, and the ALEPH group has used the CP -odd tensor observable T_{33} , which can give information only on the real part of the τ WDM. Specifically the upper limit on the real and imaginary parts of the τ WDM obtained at OPAL are $|Re(d_Z)| \leq 7.8 \times 10^{-18} \text{ecm}$ and $|Im(d_Z)| \leq 4.5 \times 10^{-17} \text{ecm}$ with 95% confidence level, while the upper limit on the real part of the τ WDM obtained at ALEPH is $|Re(d_Z)| \leq 1.5 \times 10^{-17} \text{ecm}$.

Compared with the present constraints on the τ WDM, those on the τ EDM are rather weak and, in contrast to the OPAL and ALEPH measurements, the reported measurements[36] have used indirect methods based on the CP -even observables such as the differential cross section of $e^+e^- \rightarrow \tau^+\tau^-$ [52], the partial decay widths of $Z \rightarrow \tau^+\tau^-$ [53] and $Z \rightarrow \tau^+\tau^-\gamma$ [54]. We claim that *the measurements may not be regarded as genuine τ EDM measurements and therefore the quoted values have to be replaced by those by direct measurements through CP -odd observables*[55]. Surprisingly, no direct τ EDM measurements have been reported in the literature. Therefore, we strongly suggest the τ EDM to be measured directly at the existing collider facilities TRISTAN, LEP II and CLEO II.

In the present report, first of all, we present a rough comparison of the potential of TRISTAN and LEP II in the τ EDM measurements based on the following experimental parameters:

$$\begin{aligned} \text{TRISTAN} &: \sqrt{s} = 60 \text{GeV}, \quad \mathcal{L}_{ee} = 271 \text{pb}^{-1}, \\ \text{LEP II} &: \sqrt{s} = 180 \text{GeV}, \quad \mathcal{L}_{ee} = 500 \text{pb}^{-1} \end{aligned} \quad (28)$$

In the above quoted values, we note that TRISTAN has already accumulated the integrated luminosity 271 pb^{-1} [56], but LEP II is assumed to eventually accumulate 500 pb^{-1} , which is the designed integrated luminosity per year.

For the sake of simplicity, we consider the simplest semileptonic decay of the τ lepton, $\tau \rightarrow \pi \nu_\tau$ whose branching ratio is 11%. Certainly, to obtain better constraints, all the reconstructable decay channels of the τ should be included. Given the branching fraction (11%), TRISTAN and LEP II yield ninety-seven and twenty-three events of the sequential process $e^+e^- \rightarrow \tau^+\tau^- \rightarrow (\pi^+\bar{\nu}_\tau)(\pi^-\nu_\tau)$. To make a rough estimate of the sensitivities to be obtained, we use the same set of CP -odd observables as used in the top-quark pair production at NLC in Section 2, that is to say, the CPT -even A_1 and T_{33} and the CPT -odd A_E , A_2 , and Q_{33} . All the CP -odd observables are constructed from the momenta of the electron in the initial state and two pions in the final state.

We exhibit in Fig. 8 the $1\text{-}\sigma$ allowed region of the CP -violation parameters c_γ and c_Z , which are numerically related with the τ EDM and WDM as follows

$$d_\tau^{\gamma,Z} = 1.1 \times 10^{-14} c_{\gamma,Z} (\text{ecm}). \quad (29)$$

Incidentally, it is clear that TRIATAN and LEP II can not compete with LEPI in measuring the τ WDM as can be seen clearly by comparing the bound in the figure and the values quoted in the first paragraph of the present section. So, we take into account the




Figure 8: The $1\text{-}\sigma$ allowed region (left part) of $Re(c_\gamma)$ and $Re(c_Z)$ through A_1 (solid) and T_{33} (long-dashed), and that (right part) of $Im(c_\gamma)$ and $Im(c_Z)$ through A_E (solid) and A_2 (long-dashed) at TRISTAN and LEP II with the experimental parameter values (28). The lines for Q_{33} are overlapped with those for A_E .

LEPI measurements and then we derive the $1\text{-}\sigma$ allowed range for the τ EDM by taking the x -axis cut values. Before listing the numerical values, we note that the CPT -odd observables A_E and Q_{33} provide very similar constraints on the imaginary part of the τ EDM so that their $1\text{-}\sigma$ boundary lines can be hardly distinguished in Fig. 8. Combined with the LEPI results, the $1\text{-}\sigma$ sensitivities to the τ EDM are

$$\begin{aligned} \text{TRISTAN} &: |Re(d_\tau^\gamma)| \leq 5.3 \times 10^{-16}(\text{ecm}), \quad |Im(d_\tau^\gamma)| \leq 1.1 \times 10^{-15}(\text{ecm}), \\ \text{LEPII} &: |Re(d_\tau^\gamma)| \leq 4.4 \times 10^{-16}(\text{ecm}), \quad |Im(d_\tau^\gamma)| \leq 2.8 \times 10^{-15}(\text{ecm}), \end{aligned} \quad (30)$$

TRISTAN and LEP II can measure the real part of the τ EDM with similar sensitivities, which may become comparable with those quoted by the PDG group, in the case that all the reconstructable decay modes of the tau lepton are included.

On the other hand, the e^+e^- storage ring CESR has accumulated the integrated luminosity of about 3.5 fb^{-1} at the c.m. energy $\sqrt{s} = 10.6 \text{ GeV}$ in the years between 1990 and 1994[57]. It corresponds to the production of about 3.02×10^6 τ pairs, which is about four-hundred times larger than the number of τ pairs at TRISTAN with its integrated luminosity 271 pb^{-1} . Therefore, a great improvement of the sensitivities to the τ EDM is expected. As a matter of fact, as shown in Fig. 9, the $1\text{-}\sigma$ allowed range for $Re(c_\gamma)$ is about four times smaller than that at TRISTAN and LEP II, while the $1\text{-}\sigma$ allowed range of $Im(c_\gamma)$ is about fifteen times smaller than that at TRISTAN. Quantitatively, the $1\text{-}\sigma$

sensitivities to the τ EDM, if the LEP1 results on the τ WDM are included, are

$$|Re(d_\tau^\gamma)| \leq 1.5 \times 10^{-16}(ecm), \quad |Im(d_\tau^\gamma)| \leq 6.7 \times 10^{-17}(ecm). \quad (31)$$




Figure 9: (a) The 1- σ allowed region of $Re(c_\gamma)$ and $Re(c_Z)$ through A_1 (solid) and T_{33} (long-dashed) and (b) that of $Im(c_\gamma)$ and $Im(c_Z)$ through A_E (solid) and Q_{33} (dot-dashed) with the 3.02×10^6 τ pairs at the CLEOII energy $\sqrt{s} = 10.6$ GeV.

The quoted indirect limits of the absolute value of the τ EDM is $5 \times 10^{-17}(ecm)$, which is still three times smaller than the expected limits from the CLEOII measurements. However, we should note that only the simplest decay, $\tau \rightarrow \pi\nu_\tau$, whose branching fraction is 11%, has been considered in our analysis. Therefore, we expect that including other dominant decay modes such as $\tau \rightarrow \rho\nu_\tau, a_1\nu_\tau, \dots$ enable us to easily improve the sensitivities more than five times. The more detailed work on the CLEOII measurements of the τ EDM with all the reconstructable decay modes included is in progress[58].

3.2 CP Violation in Semileptonic τ Decays

An observation of CP violation in τ decays requires the existence of not only a CP violating phase but also the interference of processes with different CP phases. One can have in general a CP violating phase between the W -exchange diagram and the charged-scalar-exchange diagram in models beyond the SM such as MHD and scalar leptoquark (SLQ) models. On the other hand, two resonance states with large width-to-mass ratios[36], which decay to the same final states, leads to an interference of the two processes with different CP phases.

The decay amplitudes of $\tau \rightarrow 3\pi\nu_\tau$ and $\tau \rightarrow K\pi\nu_\tau$ have contributions from the two overlapping resonances (a_1, π') and (K^*, K_0^*) with different spins and relatively large width-to-mass ratios. Here we should note that the parameters of the resonances are not so accurately determined. In the τ decays, various phenomenological parameterizations[59, 60, 61, 62] of the form factors have been employed to analyze experimental data. Keeping in mind the uncertainty of the resonance parameters, we simply adopt the parameterization of the τ -decay library TAUOLA[59] for the masses and widths of the resonances tabulated as follows

resonances	a_1	π'	K^*	K_0^*
J^P	1^+	0^-	1^-	0^+
mass (GeV)	1.25	1.30	0.89	1.43
width (GeV)	0.6	0.3	0.05	0.29

The semileptonic decay modes of the τ are promising for the detection of CP violation for the following reasons. First, no tagging of the other τ is necessary. Second, the decay modes can be measured not only at the CESR and CERN e^+e^- Collider LEP but also at the planned B factories and τ -charm factories[17] where many τ leptons (10^7 to 10^8) are expected to be produced.

There is, however, an experimental difficulty; Since the neutrinos escape detection, it is in general very difficult to reconstruct the τ rest frame. Nevertheless, there are two situations where the τ rest frame can be actually reconstructed. One is τ -pair production close to threshold where τ leptons are produced at rest. This possibility can be realized at future τ -charm factories. The other is when both τ leptons decay semileptonically. In the latter case impact parameter methods allow us to reconstruct the τ rest frame even for τ 's in flight. Obviously, the impact parameter method[63] requires the full identification of the decay product momenta. Therefore, we use the decay modes $\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau$ in the 3π mode and $\tau^- \rightarrow (K_s^0 \rightarrow \pi^+ \pi^-) \pi^- \nu_\tau$ in the $K\pi$ mode, of which the branching fractions are 6.8% and 0.33%, respectively.

Generally, the matrix element for the semileptonic τ decays can be cast into the form

$$M = \sqrt{2}G_F \left[(1 + \chi) \bar{u}(k, -) \gamma^\mu P_- u(p, \sigma) J_\mu + \eta \bar{u}(k, -) P_+ u(p, \sigma) J_S \right], \quad (32)$$

where G_F is the Fermi constant, p and k are the four momenta of the τ lepton and the tau neutrino, respectively, and η is a parameter determining the size of the scalar contribution. J_μ and J_S are the vector hadronic current and the scalar hadronic current given by

$$\begin{aligned} J_\mu^{3\pi} &= \cos \theta_C \langle 3\pi | \bar{d} \gamma_\mu (1 - \gamma_5) u | 0 \rangle, & J_S^{3\pi} &= \cos \theta_C \langle 3\pi | \bar{d} (1 + \gamma_5) u | 0 \rangle, \\ J_\mu^{K\pi} &= \sin \theta_C \langle (K\pi)^- | \bar{s} \gamma_\mu u | 0 \rangle, & J_S^{K\pi} &= \sin \theta_C \langle (K\pi)^- | \bar{s} u | 0 \rangle, \end{aligned} \quad (33)$$

where θ_C is the Cabibbo angle. The explicit form of J_μ is found in the τ decay library TAUOLA and the hadronic scalar current J_S can be determined by the Dirac equation from the J_μ current.

We can now construct a CP -even sum Σ and a CP -odd difference Δ of the differential $\tau^\pm$ decay rates:

$$\begin{aligned} \Sigma &= \frac{d\Gamma}{d\Phi_{X^-}} [\tau^- \rightarrow X^- \nu_\tau] + \frac{d\Gamma}{d\Phi_{X^+}} [\tau^+ \rightarrow X^+ \bar{\nu}_\tau], \\ \Delta &= \frac{d\Gamma}{d\Phi_{X^-}} [\tau^- \rightarrow X^- \nu_\tau] - \frac{d\Gamma}{d\Phi_{X^+}} [\tau^+ \rightarrow X^+ \bar{\nu}_\tau] \}, \end{aligned} \quad (34)$$

where $X = 3\pi, K\pi$ and $\Phi_{X^\pm}$ are the full kinematic variables of the detectable final states X^+ and X^- , which should be related with each other under CP transformations. Then,

the CP -odd Δ is proportional to the imaginary part of the parameter ξ , which is given by

$$\xi_{3\pi} = \frac{m_{\pi'}^2}{(m_u + m_d)m_\tau} \left(\frac{\eta}{1 + \chi} \right), \quad \xi_{K\pi} = \frac{m_{K_0^*}^2}{(m_s - m_d)m_\tau} \left(\frac{\eta}{1 + \chi} \right). \quad (35)$$

We re-emphasize that every CP asymmetry requires not only a non-vanishing $Im(\xi)$ but also the interference between the longitudinal mode of the vector meson and the scalar meson and the interference is proportional to the decay constants $f_{\pi'}$ and $f_{K_0^*}$. The value of the π' decay constant, $f_{\pi'} = 0.02\text{--}0.08$ GeV estimated in Ref. [61] and $f_{\pi'} = 0.02$ GeV quoted in TAUOLA are quite likely invalid because the mixing between the chiral pion field and a massive pseudoscalar $q\bar{q}$ bound state should be considered. So, we have reconsidered $f_{\pi'}$ in the chiral Lagrangian framework and have shown that it actually vanishes in the chiral limit due to the mixing of π' with the chiral pion field[14]. $f_{\pi'}$ is indeed proportional to the square of the pion mass m_π and so it may be much smaller than the value quoted in TAUOLA. This small $f_{\pi'}$ value is also supported by arguments from QCD sum rules. On the other hand, $f_{K_0^*}$ can be well determined phenomenologically by using the well-measured K_0^* width and the QCD sum rule technique. Considering possible uncertainties in our estimates we use in our numerical analysis $f_{\pi'} = (1 \sim 5) \times 10^{-3}$ GeV for the π' decay constant, which is a slightly broader range than estimated, and $f_{K_0^*} = 45$ MeV for the K_0^* decay constant, which is rather well estimated.

As possible new sources of CP violation detectable in the τ decays we consider new scalar-fermion interactions which preserve the symmetries of the SM. Under those conditions there exist only four types of scalar-exchange models[64], which can contribute to the decays, $\tau \rightarrow 3\pi\nu_\tau$ and $\tau \rightarrow (K\pi)\nu_\tau$. One is the MHD model[42] and the other three are scalar-leptoquark (SLQ) models[64, 65]. The quantum numbers of the three leptoquarks under the gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$ are

$$\begin{aligned} \Phi_1 &= \left(3, 3, \frac{7}{6} \right) \quad (\text{model I}), \\ \Phi_2 &= \left(3, 1, -\frac{1}{3} \right) \quad (\text{model II}), \\ \Phi_3 &= \left(3, 3, -\frac{1}{3} \right) \quad (\text{model III}), \end{aligned} \quad (36)$$

respectively. The hypercharge Y is defined to be $Q = I_3 + Y$.

The constraints on the CP -violation parameters $\xi_{3\pi}$ and $\xi_{K\pi}$ depend upon the values of the u , d and s current quark masses, which are taken to be

$$m_u = 5 \text{ MeV}, \quad m_d = 9 \text{ MeV}, \quad m_s = 320 \text{ MeV}. \quad (37)$$

The present experimental constraints on the CP -violation parameters in the MHD model, which have been extensively reviewed in Ref. [42], are

$$\frac{\Gamma(B \rightarrow X\tau\nu_\tau)}{\Gamma(K^+ \rightarrow \pi^+\nu\bar{\nu})} \Rightarrow |Im(\xi_{MHD})|_{3\pi} < 0.28, \quad |Im(\xi_{MHD})|_{K\pi} < 0.48, \quad (38)$$

for the charged Higgs mass $M_H = 45$ GeV. There are at present no direct constraints on the CP -violating parameters in the SLQ models. Assuming that the three leptoquark couplings to fermions are universal, we can roughly estimate the constraints on the SLQ CP -violating parameters. The constraints[36, 66, 67] on the CP -violation parameters from various low energy experiments then are

$$\begin{aligned} \Gamma(K_L \rightarrow \mu e) &\Rightarrow |Im(\xi_{SLQ}^I)|_{3\pi} < 1.5 \times 10^{-3}, \\ \frac{\Gamma(\mu\text{Ti} \rightarrow e\text{Ti})}{\Gamma(\mu\text{Ti} \rightarrow \text{capture})} &\Rightarrow |Im(\xi_{SLQ}^{II})|_{3\pi} < 0.9 \times 10^{-3}, \end{aligned} \quad (39)$$

$$\begin{aligned} D\bar{D} \text{ mixing} &\Rightarrow \begin{cases} |Im(\xi_{SLQ}^I)|_{K\pi} < 4 \times 10^{-2}, \\ |Im(\xi_{SLQ}^{II})|_{K\pi} < 6 \times 10^{-2}, \end{cases} \\ \Gamma(K \rightarrow \pi\nu\bar{\nu}) &\Rightarrow |Im(\xi_{SLQ}^{III})|_{K\pi} < 10^{-5}, \end{aligned} \quad (40)$$

Compared to the constraint on the $Im(\xi_{MHD})$, the constraints on the SLQ CP -violation parameters are much more severe. Especially, the constraint on $Im(\xi_{SLQ}^{III})$ is so severe that no CP -violation effects are expected to be observed. Therefore we will not consider this type-III SLQ model any longer.

The following numerical analysis is made for the maximally allowed values of the CP -violation parameters and the number of τ leptons required to detect CP violation is estimated by using the optimal CP -odd asymmetry[21]

$$w_{\text{opt}} = \frac{\Delta}{\Sigma}. \quad (41)$$

Table 5: The maximal expected size of ε_{opt} and the number of τ leptons, N , for detection with the ε_{opt} at the 2σ level in the 3π and $K\pi$ modes for $f_{\pi'} = 1 \sim 5$ MeV and $f_{K_0^*} = 45$ MeV with the maximally-allowed values for the CP -violation parameters.

Model	$\varepsilon_{\text{opt}}^{3\pi}(\%)$	$\varepsilon_{\text{opt}}^{K\pi}(\%)$	$N^{3\pi}$	$N^{K\pi}$
MHD	$0.13 \sim 0.67$	2.3	$(0.13 \sim 3.3) \times 10^7$	2.3×10^6
SLQI	$(0.7 \sim 3.6) \times 10^{-3}$	0.2	$(0.5 \sim 11) \times 10^{11}$	3.4×10^8
SLQII	$(0.4 \sim 2.2) \times 10^{-3}$	0.3	$(1.2 \sim 33) \times 10^{11}$	1.5×10^8

Table 5 shows the expected size of ε_{opt} , along with the number of τ leptons, N , required to obtain the $2\text{-}\sigma$ signal with the optimal asymmetry, ε_{opt} , in the MHD model and the two SLQ models for the CP -violation parameter values. The values of ε_{opt} and the corresponding N values in Table 5 show that the CP -violating effects from the MHD model can be detected with less than 10^7 τ leptons, while still more than 10^8 τ leptons are required to see CP violation in the SLQ models. In light of the fact that about 10^7 and 10^8 τ leptons are produced yearly at B and τ -Charm factories, respectively, CP violation from the MHD model can hopefully be observed.

4 Summary and Conclusion

Let us summarize the results presented in the present report.

First of all, we have made a systematic study of observable experimental CP -odd distributions in the e^+e^- annihilation processes, $e^+e^- \rightarrow W^+W^-$ and $e^+e^- \rightarrow t\bar{t}$ and of initial CP -odd two-photon polarization configurations in the two-photon fusion processes, $\gamma\gamma \rightarrow W^+W^-$ and $\gamma\gamma \rightarrow t\bar{t}$ at NLC, which could serve as tests of possible anomalous CP -odd three-boson and four-boson couplings and top-quark EDM and WDM couplings.

We have found that backscattering linearly polarized Compton laser light off the electron or positron beam allows us to prepare two types of initial CP -odd two-photon polarization configurations and that the sensitivities through the two CP -odd asymmetries to the CP -violation parameters depend strongly on the parameter x , i.e. the laser beam frequency. So, from an experimental point of view, it is very crucial to be able to adjust the initial laser beam frequency in probing CP violation.

Several works have shown that new models beyond the SM such as MHD and MSSM could have CP -odd parameters of the size[68] which are close to the limits obtainable from W -pair production both in e^+e^- annihilation and two-photon fusion at an NLC. On the other hand, the expected size of the top-quark EDM and WDM is smaller than the reachable limits by an order of magnitude. However, it is expected that the two-photon mode with higher c.m. energies and adjustable laser frequencies could approach very close to the expectation.

Secondly, we have pointed out that no direct measurements of the τ EDM have been reported, while two very good measurements of the τ WDM have been performed by OPAL[50] and ALEPH[51]. For that reason, we have suggested any direct measurements of the τ EDM to be immediately presented at the presently existing collider facilities, TRISTAN, LEP II and CLEO II. We have compared the potential of TRISTAN, LEP II and CLEO II in the τ EDM measurements by using the simplest τ decay mode $\tau \rightarrow \pi\nu_\tau$. As a result, we have found that, when all the constructible τ decay modes are included, CLEO II should enable us to obtain limits similar to or better than the indirect limits quoted in the literature[36].

Thirdly, the semileptonic τ decays, $\tau \rightarrow 3\pi\nu_\tau$ and $\tau \rightarrow K\pi\nu_\tau$, involve two different intermediate resonances with large ratios of widths to masses so that we have proposed to investigate them as very promising decay channels to probe CP violation which stems from new sources completely different from the SM CP -violation mechanism. Quantitatively, it turns out that the CP -violating effects due to the charged Higgs exchanges in the MHD model might be detected at the planned B and proposed τ -charm factories, while those in the SLQ models are hardly detectable.

Consequently, we conclude that (i) CP violation from new physics can constitute an interesting research programme at NLC, (ii) any direct τ EDM measurements should be made immediately at the presently existing collider experiments, in particular at CLEO II to replace the indirect limits quoted in the literature, and (iii) the semileptonic τ decays are worthwhile to be investigated as promising channels to probe CP violation from new physics. Any CP -violation phenomenon in the processes under consideration, if

discovered, implies new CP -violation mechanism, which can have far-reaching physical consequences.

Acknowledgments

The authors would like to thank Profs. O. Nachtmann and M. Tanabashi for useful comments and fruitful discussions. SYC thank the KEK Theory Group and the DESY theory group for their kind hospitality during his stay. The work of SYC was supported in part by the KOSEF-DFG large collaboration project, Project No. 96-0702-01-01-2 and in part by the Korean Science and Engineering Foundation (KOSEF) and Korean Federation of Science and Technology Societies through the Brain Pool program and the work of MSB was supported in part by KOSEF and Research University Fund of College of Science at Yonsei university supported by the Ministry of Education, Korea.

References

- [1] S.L. Glashow, Nucl. Phys. **22** (1961) 579; S. Weinberg, Phys. Rev. Lett. **19** (1967) 1264; A. Salam, in *Elementary Particle Theory*, ed. N. Svartholm (Almqvist and Wiksells, Stockholm, 1969), p. 367.
- [2] H. Georgi, Ann. Rev. Nucl. Part. **43** (1994) 209; Nucl. Phys. **B361** (1991) 339.
- [3] J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay, Phys. Rev. Lett. **13** (1964) 138.
- [4] N. Cabibbo, Phys. Rev. Lett. **10** (1963) 531.
- [5] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49** (1973) 652.
- [6] L. Wolfenstein, Phys. Rev. Lett. **13** (1964) 562.
- [7] E. D. Kolb and M. S. Turner, *The Early Universe* (Addison-Wesley-Publishing, New York, 1990).
- [8] A. D. Sakharov, ZhETF Pis. Red. **5** (1967) 32; JETP Lett. **5** (1967) 24.
- [9] G.R. Farrar and M.E. Shaposhnikov, Phys. Rev. D**50** (1994) 774; M.B. Gavela *et al.*, Nucl. Phys. **B430** (1994) 382; P. Huet and E. Sather, Phys. Rev. D**51** (1995) 379.
- [10] E. P. Shabalin, Sov. J. Nucl. Phys. **28** (1978) 75; J. Donoghue, Phys. Rev. D**18**, (1978) 1632; I.B. Khriplovich, Sov. J. Nucl. Phys. **44** (1986) 659; Phys. Lett. **173B** (1986) 193; A. Czarnecki and B. Krause, Acta. Phys. Pol. B **28** (1997) 829; Phys. Rev. Lett. **78** (1997) 4339.
- [11] F. Hoogeveen, Nucl. Phys. **B341** (1990) 322.

- [12] S.Y. Choi, K. Hagiwara, and M.S. Baek, Phys. Rev. D**54** (1996) 6703; M.S. Baek and S.Y. Choi, Preprint No. YUMS 97-5, SNUTP 97-028.
- [13] S.Y. Choi and K. Hagiwara, Phys. Lett. **359B** (1995) 369; M.S. Baek, S.Y. Choi and C.S. Kim, Preprint No. YUMS 97-7, SNUTP 97-035, hep-ph/9704312.
- [14] S.Y. Choi, K. Hagiwara and M. Tanabashi, Phys. Rev. D**52** (1995) 1614.
- [15] Proceedings of the 1st International Workshop on “Physics and Experiments with Linear e^+e^- Colliders” (Saariselka, Finland, September 1991), eds. R. Orava, P. Eerola, and M. Nordberg (World Scientific, Singapore, 1992); Proceedings of the 2nd International Workshop on “Physics and Experiments with Linear e^+e^- Colliders” (Waikoloa, Hawaii, April 1993), eds. F.A. Harris, S.L. Olsen, S. Pakvasa, and X. Tata (World Scientific, Singapore, 1993), Proceedings of the 3rd International Workshop on “Physics and Experiments with Linear e^+e^- Colliders” (Iwate, Japan, September 1995), eds. A. Miyamoto, Y. Fujii, T. Matsui, and S. Iwata, (World Scientific, Singapore, 1996).
- [16] I.F. Ginzburg, G.L. Kotkin, V.G. Serbo, S.L. Panfil and V.I. Telnov, Sov. ZhETF Pis'ma **34**, 514 (1981) [JETP Lett. **34**, 491 (1982)]; Nucl. Instr. and Meth. **205**, 47 (1983); I.F. Ginzburg, G.L. Kotkin, S.L. Panfil, V.G. Serbo, and V.I. Telnov, *ibid* **219**, 5 (1984); V.I. Telnov, *ibid* **A294**, 72 (1992); J.F. Gunion and H.E. Haber, Phys. Rev. D **48** (1993) 5109; S.Y. Choi and F. Schrempp, Phys. Lett. **272B**, 149 (1991); E. Yehudai, Phys. Rev. D**44**, 3434 (1991); Ph. D Thesis, SLAC-PUB-383.
- [17] J.J. Gomez-Cadenas, in *Proceedings of the Third Workshop on the Tau-Charm Factory*, Marbella, Spain, 1993, edited by J. Kirkby and R. Kirkby (Editions Frontiers, Gif-sur-Yvette, France, 1994); B.C. Barish and R. Stroynowski, Phys. Rep. **157** (1988) 1; K. Riles, Int. J. Mod. Phys. **A7** (1992) 7647; M.L. Perl, Rep. Prog. Phys. **55** (1992) 653.
- [18] W. Bernreuther and A. Brandenburg, Phys. Lett. **314B** (1993) 104; W. Bernreuther, J. P. Ma, and B. H. J. McKellar, Phys. Rev. D**51** (1995) 2475; H. Anlauf, W. Bernreuther, and A. Brandenburg, Phys. Rev. D**52** (1995) 3803 and references therein.
- [19] F. Cuypers and S.D. Rindani, Phys. Lett. **343B** (1995) 333.
- [20] D. Chang, W. Y. Keung, and I. Phillips, Nucl. Phys. **B408** (1993) 286; **429** (1994) 255 (E).
- [21] D. Atwood and A. Soni, Phys. Rev. D**45** (1992) 2405.
- [22] P. Poulose and S.D. Rindani, Phys. Lett. **349B** (1995) 379.
- [23] K. Hagiwara, R.D. Peccei, D. Zeppenfeld and K. Hikasa, Nucl. Phys. **B282** (1987) 253.

- [24] G. Gounaris, D. Schildknecht, and F.M. Renard, Phys. Lett. **263B**, 291 (1991); M.B. Gavela, F. Iddir, A. Le Yaouanc, L. Oliver, O. Pène, and J.C. Raynal, Phys. Rev. D**39**, 1870 (1989); A. Bilal, E. Massó, and A. De Rújula, Nucl. Phys. **B355**, 549 (1991).
- [25] P. Kalyniak, P.Madsen, N. Sinha, and R. Sinha, Phys. Rev. D**52**, 3826 (1995); D. Chang, W. Y. Keung, and I. Phillips, Phys. Rev. D**48** (1993) 4045; G. Gounaris, D. Schildknecht, and F. M. Renard, Phys. Lett. **263B** (1991) 291; M. B. Gavela, F. Iddir, A. Le Yaouanc, L. Oliver, O. Pène, and J. C. Raynal, Phys. Rev. D**39** (1989) 1870; A. Bilal, E. Massó, and A. De Rújula, Nucl. Phys. **B355** (1991) 549.
- [26] G.J. Gounaris and G.P. Tsirgoti, Preprint No. THES-TP 97/03, E-preprint No. hep-ph/9703446.
- [27] D. Atwood, G. Eilam, and A. Soni, Phys. Rev. Lett. **71** (1993) 492; B.Grzadkowski and J. F. Gunion, Phys. Lett. **287B** (1992) 237.
- [28] J. Wudka, Nucl. Phys. Proc. Suppl. 37A (1994) 211.
- [29] G. Bélanger and G. Couture, Phys. Rev. D**49** (1994) 5720.
- [30] G.L. Kane, G.A. Ladinsky, and C.-P. Yuan, Phys. Rev. D**45** (1992) 124; C.R. Schmidt and M.E. Peskin, Phys. Rev. Lett. **69** (1992) 410; D. Atwood, A. Aepli, and A. Soni, Phys. Rev. Lett. **69** (1992) 2754.
- [31] K. Hikasa, Phys. Lett. **143B**, 266 (1984); Phys. Rev. D**33**, 3203 (1986).
- [32] M. Krämer, J. Kühn, M.L. Stong and P.M. Zerwas, Z. Phys. **C64** (1994) 21.
- [33] W. Buchmüller and D. Wyler, Nucl. Phys. **B268** (1986) 621; C. J. C. Burges and H. J. Schnitzer, *ibid* **B228** (1983) 464; C. N. Leung, S. T. Love and S. Rao, Z. Phys. **C31** (1986) 433; K. Hagiwara, S. Ishihara, R. Szalapski, D. Zeppenfeld, Phys. Rev. D**48** (1993) 2182; K. Hagiwara and M. Stong, Z. Phys. **C62** (1994) 99.
- [34] W. Bernreuther and O. Nachtmann, Phys. Rev. Lett. **63** (1989) 2787; W. Bernreuther, O. Nachtmann, P. Overmann and T. Schröder, Nucl. Phys. **B388** (1992) 53.
- [35] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **74** (1995) 2626; S. Abachi *et al.* (D0 Collaboration), *ibid.* **74** (1995) 2632.
- [36] Particle Data Group, L. Montanet *et al.*, Phys. Rev. D**54** (1996) 94.
- [37] Ikaros I. Y. Bigi, Yu L. Dokshitser, V. Khoze, J. Kühn and P. Zerwas, Phys. Lett. **181B** (1986) 157.
- [38] S.M. Barr and W.J. Marciano, in *CP Violation*, ed. C. Jarlskog (World Scientific, Singapore, 1990).

- [39] A. Soni and R.M. Xu, Phys. Rev. Lett. **69** (1992) 33.
- [40] M. Jezabek and J.H. Kühn, Nucl. Phys. **B320** (1989) 20; A. Czarnecki, M. Jezabek and J.H. Kühn, Nucl. Phys. **B351** (1991) 70.
- [41] Y.-S. Tsai, Phys. Rev. D **4** (1971) 2821; S.Y. Pi and A.I. Sanda, Ann. Phys. (N.Y.) **106** (1977) 16; H. Kühn and F. Wagner, Nucl. Phys. **B236** (1984) 16; K. Hagiwara, A.D. Martin and D. Zeppenfeld, Phys. Lett. **235B** (1989) 198; B.K. Bullock, K. Hagiwara and A.D. Martin, Phys. Rev. Lett. **67** (1991) 3055; Nucl. Phys. **B395** (1993) 499; P. Privitera, Phys. Lett. **308B** (1993) 163.
- [42] Y. Grossman, Nucl. Phys. **B426** (1994) 355 and references therein.
- [43] D. Atwood, G. Eilam and A. Soni, Phys. Rev. Lett. **71** (1993) 492; A.F. Falk, Z. Ligeti, M. Neubert and Y. Nir, Phys. Lett. **326B** (1994) 145; Y. Grossman and Z. Ligeti, *ibid.* **332**, (1994) 373.
- [44] W. Bernreuther, O. Nachtmann and P. Overmann, Z. Phys. **C48** (1993) 78 and references therein.
- [45] U. Kilan, J.G. Körner, K. Schilcher and W.L. Wu, Z. Phys. **C62** (1994) 413.
- [46] B. Ananthanarayan and S.D. Rindani, Phys. Rev. Lett. **73** (1994) 1215.
- [47] Y.S. Tsai, Phys. Lett. **378B** (1996) 272 and references therein.
- [48] T. Huang, W. Lu and Z. Tao, Phys. Rev. D **55** (1997) 1643.
- [49] J.H. Kühn and E. Mirkes, Phys. Lett. **398B** (1997) 407.
- [50] R. Akers *et al.* (OPAL Collaboration), Z. Phys. **C66** (1995) 31.
- [51] D. Buskulic *et al.* (ALEPH Collaboration), Phys. Lett. **346B** (1995) 371.
- [52] F. del Aguila and M. Sher, Phys. Lett. **252B** (1990) 116.
- [53] J.A. Grifols and A. Mendez, Phys. Lett. **255B** (1991) 611.
- [54] R. Escribano and E. Masso, Phys. Lett. **301B** (1993) 419.
- [55] O. Nachtmann, private communication.
- [56] H. Hanai *et al.* (Venus Collaboration), Preprint No. KEK preprint 96-171 (1996).
- [57] J.P. Alexander *et al.* (CLEO Collaboration), Preprint No. CLNS 97/1480, CLEO 97-9.
- [58] M.S. Baek, S.Y. Choi and K. Hagiwara, in preparation.

- [59] S. Jadach, J.H. Kühn and Z. Was, *Comput. Phys. Commun.* **64** (1991) 275; M. Jezabek, Z. Was, S. Jadach, and J.H. Kühn, *ibid.* **70** (1992) 69; S. Jadach, Z. Was, R. Decker, and J.H. Kühn, *ibid.* **76** (1993) 361.
- [60] N.A. Tornqvist, *Z. Phys.* **CC36** (1987) 695; **40** (1988) 632; M.G. Bowler, *Phys. Lett.* **209B** (1988) 99; Yu.P. Ivanov, A.A. Osipov and M.K. Volkov, *Z. Phys.* **C49** (1991) 563.
- [61] N. Isgur, C. Morningstar and C. Reader, *Phys. Rev. D* **39** (1989) 1357.
- [62] R. Decker, E. Mirkes, R. Sauer and Z. Was, *Z. Phys.* **C58** (1993) 445.
- [63] J.H. Kühn, *Phys. Lett.* **313B** (1993) 458.
- [64] A.J. Davies and X.-G. He, *Phys. Rev. D* **43** (1991) 225.
- [65] L.J. Hall and L.J. Randall, *Nucl. Phys.* **B274** (1986) 157; J.F. Nieves, *ibid.* **B189** (1981) 382; W. Buchmüller, R. Rückl and D. Wyler, *Phys. Lett.* **191B** (1987) 44.
- [66] C. Dohmen et al., *Phys. Lett.* **317B** (1993) 631.
- [67] W.J. Marciano and A. Sirlin, *Phys. Rev. Lett.* **71** (1994) 3629.
- [68] W. Marciano and A. Queijeiro, *Phys. Rev. D* **33**, 3449 (1986); F. Boudjema, K. Hagiwara, C. Hamzaoui, and K. Numata, *ibid D* **43**, 2223 (1991); A. De Rijula, M. Gavela, O. Pène, and F. Vegas, *Nucl. Phys.* **B357**, 311 (1991); see also S. Barr and W. Marciano, *CP Violation*, ed. C. Jarlskog (World Scientific, Singapore, 1989); D. Chang, W.Y. Keung and J. Liu, *Nucl. Phys.* **B355** (1991) 295; R. López-Mobilia and T.H. West, *Phys. Rev. D* **51** (1995) 6495.